

Mat 1160
WEEK 6

Dr. N. Van Cleave

Spring 2010

Student Responsibilities – Week 6

- ▶ **Reading:**

 - This week: Textbook, Section 2.5: Infinite Sets

 - Next week: Textbook, Sections 3.1–3.2: Logic and Truth Tables

- ▶ Summarize Sections & Work Examples
Sample Sec 1.1 and 1.2 are on our web site
- ▶ Attendance
- ▶ Recommended exercises:
 - ▶ Section 2.5: (all 1–6), evens 8–46
 - ▶ Chapter test

THE DENZEL WASHINGTON VENN DIAGRAM



Sec 2.5 Infinite Sets & Their Cardinalities



- ▶ **Cardinal Number** or **Cardinality**: of a finite set is the number of elements that it contains.

- ▶ **One-to-one (1-1) Correspondence**: the elements in two sets can be matched together in such a way that each element is paired with exactly one unique element from the other set.



One-to-one Correspondence Example

For example, we can find a 1–1 correspondence between the sets
 $\{ a, b, c, d \}$ and $\{ 1, 2, 3, 4 \}$ by pairing
(a,1), (b,2), (c,3), and (d,4).

Other matches would work also: **(1,b), (2,c), (3,d), and (4,a)**




- Examples of correspondences which are **not** 1-1:



Set Equivalence

- ▶ Two sets A and B are **equivalent**, ($A \sim B$), if they can be placed in a **1–1 correspondence**.
- ▶ \aleph_0 , **Aleph-naught** or **Aleph-null**, is the **cardinality** of the natural (or counting) numbers, $\{1, 2, 3, \dots\}$, a **countably infinite** set.
- ▶ If we can show a 1–1 correspondence between some set, A , and the natural numbers, we say that A also has cardinality \aleph_0 .
- ▶ **Thus, to show a set A has cardinality \aleph_0 , we need to find a 1–1 correspondence between A and the set of natural numbers, \mathbb{N}**

► Can we show $|\mathbb{W}| = \aleph_0$? **Yes:**




► Notice that we just proved that $\aleph_0 = 1 + \aleph_0$!

That is just too weird!


Infinite Sets

- ▶ A set is **infinite** if it can be placed in a one-to-one correspondence with a **proper** subset of **itself**.
- ▶ Can we show that the set of integers has cardinality \aleph_0 ? Yes:



- ▶ **Good Grief!** We just showed that $\aleph_0 = 2 \times \aleph_0$! The cardinality of the counting numbers is the same as the cardinality of the integers!
- ▶ There are just as many counting numbers as there are integers. . .

We can even show that the set of **rational numbers** has cardinality \aleph_0 .



The correspondence begins. . .

$$\frac{0}{1} \leftrightarrow 1 \qquad \frac{1}{1} \leftrightarrow 2$$

$$\frac{2}{1} \leftrightarrow 3 \qquad \frac{1}{2} \leftrightarrow 4$$

$$\frac{1}{3} \leftrightarrow 5 \qquad \frac{1}{4} \leftrightarrow 6$$

- ▶ Numbers that are shaded are omitted since they can be reduced to lower terms, and were thus included earlier in the listing. (This is actually optional).
- ▶ The mapping of the positive Rational numbers to the natural numbers is a 1–1 correspondence, so it shows that the **positive Rational numbers have cardinality \aleph_0** .
- ▶ By using the method in the example for the integers, we can extend this correspondence to include negative rational numbers.
- ▶ Thus, **the set of rational numbers, \mathbb{Q} , has cardinality \aleph_0** .

- ▶ A set is **countable** if it is **finite** or if it has **cardinality** \aleph_0 .

- ▶ If a set is both **countable** and **infinite**, we call it **countably infinite**.

- ▶ Recall: $\mathbb{R} = \{ x \mid x \text{ is a number that can be written as a decimal} \}$
- ▶ We can show that the set of all real numbers, \mathbb{R} , does **not** have cardinality \aleph_0 (and in fact, is **larger** than \aleph_0) using a technique called **diagonalization**.

We will **assume** $|\mathbb{R}| = \aleph_0$
and
show this leads to a **contradiction**.

- ▶ Since we **assumed** \mathbb{R} is **countably infinite**, there is a **1-1 correspondence between it and \mathbb{N}** (by definition).

Thus some decimal number corresponds to the counting number 1, another to 2, and so on. Suppose:

$$1 \leftrightarrow .\mathbf{1}396875\dots$$

$$2 \leftrightarrow .4\mathbf{8}13863\dots$$

$$3 \leftrightarrow .75\mathbf{2}7790\dots$$

$$4 \leftrightarrow .394\mathbf{0}355\dots$$

$$5 \leftrightarrow \dots$$

and on and on....

Next we **construct** a decimal number which **cannot** be a part of this correspondence.

But we assumed $|\mathfrak{R}| = \aleph_0$, so **every** decimal number **must appear** somewhere in the correspondence list!

A Contradiction!

Let us construct a decimal number, D , according to the following:

1. The first decimal number in the given list has **1** as its first digit; let D start as $D = .2\dots$ Thus D **cannot be the first number in the list.**
2. The second decimal in the list above has **8** as its second digit; let $D = .29\dots$ Thus D **cannot be the second number in the list.**
3. The third decimal in the list has **2** as its third digit; let $D = .293\dots$ Thus D **cannot be the third number on the list.**
4. The fourth digit of the fourth decimal is **0**, so let $D = .2931\dots$
5. Continue building D in this manner

Now we get to ask: **Is D in the list**
(that we assumed **contained all decimals** numbers?)

NO, since every decimal number in the list differs from D in at least one position, **D cannot possibly be in the list!!** ($\Rightarrow \Leftarrow$)

Note:

1. We assumed **every** decimal number was in the list.
2. But the decimal number D is **not** in the list.

This presents us with a **Contradiction** — i.e.,
these statements **cannot both be true**.

Thus our original assumption was **incorrect**:

**It is not possible
to find a 1–1 correspondence between \mathfrak{R} and \mathfrak{N}**

- ▶ The set of real numbers \mathbb{R} has cardinality \mathfrak{C} (cardinality of the continuum), which is **larger** than \aleph_0 .
- ▶ The POWER SET of \mathbb{R} has cardinality **even greater** than \mathfrak{C} ... and it only gets worse from there.
- ▶ \mathbb{N} , \mathbb{W} , \mathbb{I} , and \mathbb{Q} all have cardinality \aleph_0
- ▶ Irrational numbers and \mathbb{R} have cardinality \mathfrak{C} .