

## Sec 3.2 Truth Tables and Equivalent Statements

**Conjunction:** Given two statements  $p$  and  $q$ , their conjunction is  $p \wedge q$ .

Conjunction Truth Table

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

## Conjunction Examples Determine the truth values (T/F):

- \_\_\_ Today is Tuesday **and** it is sunny.
- \_\_\_ Today is Wednesday **and** it is sunny.
- \_\_\_ The moon is made of green cheese **and** some violets are blue.
- \_\_\_ It is daytime here **and** there are **not** 1000 desks in this classroom.
- \_\_\_ This course is **MAT 1160** **and** we are learning calculus.
- \_\_\_ This course is **MAT 4870** **and** we are learning physics.
- \_\_\_  $3 < 5 \wedge 5 < 3$
- \_\_\_  $3 < 5 \wedge 5 < 8$

## Disjunction

**Disjunction.** Given two statements  $p$  and  $q$ , their (inclusive) disjunction is  $p \vee q$ .

**Inclusive disjunctions** are TRUE if either or both components are TRUE.

Disjunction Truth Table

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

## Disjunction Examples Determine the truth values:

- \_\_\_ Today is Tuesday **or** it is sunny.
- \_\_\_ Today is Wednesday **or** it is sunny.
- \_\_\_ The moon is made of green cheese **or** some violets are blue.
- \_\_\_ It is daytime here **or** there are not 100 desks in this classroom.
- \_\_\_ This course is **MAT 1160** **or** we are learning calculus.
- \_\_\_ This course is **MAT 4870** **or** we are learning physics.
- \_\_\_  $3 < 5 \vee 5 < 3$
- \_\_\_  $3 < 5 \vee 5 < 8$

## Mathematical Examples Using **or**

Statement	Reason It's True
$7 \geq 7$	$7 = 7$
$8 \geq 5$	$8 > 5$
$-7 \leq -3$	$-7 < -3$
$-3 \leq -3$	$-3 = -3$

## The Porsche & The Tiger

A prisoner must make a choice between two doors: behind one is a beautiful red Porsche, and behind the other is a hungry tiger. Each door has a sign posted on it, but only one sign is true.

**Door #1.** IN THIS ROOM THERE IS A PORSCHE AND IN THE OTHER ROOM THERE IS A TIGER.

**Door #2.** IN ONE OF THESE ROOMS THERE IS A PORSCHE AND IN ONE OF THESE ROOMS THERE IS A TIGER.

With this information, the prisoner is able to choose the correct door... Which one is it?

## Negation

**Negation.** Given a statement  $p$ , its negation is  $\sim p$ .

### Negation Truth Table

$p$	$\sim p$
<b>T</b>	<b>F</b>
<b>F</b>	<b>T</b>

## Negation Examples Determine the truth values Assume $p$ is **TRUE**, $q$ is **FALSE**, and $r$ is **FALSE**

1.  $\underline{\quad} p$
2.  $\underline{\quad} \sim p$
3.  $\underline{\quad} q$
4.  $\underline{\quad} \sim q$
5.  $\underline{\quad} r$
6.  $\underline{\quad} \sim r$
7.  $\underline{\quad} \sim p \wedge p$
8.  $\underline{\quad} p \vee \sim p$
9.  $\underline{\quad} p \wedge \sim q$
10.  $\underline{\quad} p \vee \sim q$
11.  $\underline{\quad} \sim p \wedge (q \vee \sim r)$
12.  $\underline{\quad} p \wedge (\sim q \vee r)$

## More Examples Determine the truth values

Let  $p$  represent the statement  $3 > 2$   
 $q$  represent the statement  $5 < 4$   
 $r$  represent the statement  $3 \leq 8$

1.  $\underline{\quad} p$
2.  $\underline{\quad} \sim p$
3.  $\underline{\quad} q$
4.  $\underline{\quad} \sim q$
5.  $\underline{\quad} r$
5.  $\underline{\quad} \sim r$
7.  $\underline{\quad} \sim p \wedge q$
8.  $\underline{\quad} \sim (p \wedge q)$
9.  $\underline{\quad} \sim p \vee (\sim q \vee r)$
10.  $\underline{\quad} (\sim p \wedge r) \vee (\sim q \wedge \sim p)$

## Yet More Examples

11.  $\underline{\quad}$  For some real number  $x$ ,  $x > 2$  and  $x < 8$
12.  $\underline{\quad}$  There exists a real number  $b$ ,  $b < 8$  or  $b > 2$
13.  $\underline{\quad}$  For at least one real number  $y$ ,  $y < 8$  and  $y > 12$
14.  $\underline{\quad}$  There is a real number  $m$ ,  $m < 8$  or  $m > 12$
15.  $\underline{\quad}$  For all real numbers  $x$ ,  $x < 8$  and  $x > 2$
16.  $\underline{\quad}$  For every real number  $b$ ,  $b < 8$  or  $b > 2$
17.  $\underline{\quad}$  For all real numbers  $y$ ,  $y < 8$  and  $y > 12$
18.  $\underline{\quad}$  For every real number  $m$ ,  $m < 8$  or  $m > 12$
19.  $\underline{\quad}$  For every real number  $n$ ,  $n^2 > 0$
20.  $\underline{\quad}$  For every real number  $n$ ,  $n^2 \geq 0$

## Constructing Truth Tables

Construct a Truth Table for:  $(\sim p \wedge q) \vee \sim q$

$p$	$q$	$(\sim p \wedge q) \vee \sim q$
T	T	
T	F	
F	T	
F	F	

## Construct a Truth Table for: $p \wedge (\sim p \vee \sim q)$

$p$	$q$	$p \wedge (\sim p \vee \sim q)$
T	T	
T	F	
F	T	
F	F	

### Construct the Truth Table

p	q	r	$\sim p \wedge (q \vee \sim r)$
T	T	T	
T	T	F	
T	F	T	
T	F	F	
F	T	T	
F	T	F	
F	F	T	
F	F	F	

### Construct the Truth Table

p	q	r	$(\sim p \wedge r) \vee (\sim q \wedge \sim p)$
T	T	T	
T	T	F	
T	F	T	
T	F	F	
F	T	T	
F	T	F	
F	F	T	
F	F	F	

### Some Notes of Interest

A logical statement having  $n$  component statements will have  $2^n$  rows in its truth table.

Two statements are **equivalent** if they have the same truth value in **every** possible situation.

In other words, two statements are **equivalent** if their columns in the same truth table have the same truth values.

### De Morgan's Laws

p	q	$\sim p \wedge \sim q$	$\sim (p \vee q)$
T	T		
T	F		
F	T		
F	F		

p	q	$\sim p \vee \sim q$	$\sim (p \wedge q)$
T	T		
T	F		
F	T		
F	F		