Basic Concepts of Set Theory: Symbols & Terminology

A set is a collection of objects.

A **well-defined set** has no ambiguity as to what objects are in the set or not.

For example:

- The collection of all red cars
- The collection of positive numbers
- The collection of people born before 1980
- The collection of greatest baseball players

All of these collections are sets. However, the collection of greatest baseball players is not well-defined.

Normally we restrict our attention to just well-defined sets.

word description

The set of odd counting numbers between 2 and 12

• the listing method

 $\{3,5,7,9,11\}$

• set-builder notation or defining property method

 $\{x \mid x \text{ is a counting number, } x \text{ is odd, and } x < 12\}$

Note:

- Use curly braces to designate sets,
- Use commas to separate set elements
- The variable in the set-builder notation doesn't have to be x.
- Use ellipses (...) to indicate a continuation of a pattern established before the ellipses i.e. $\{1, 2, 3, 4, ..., 100\}$
- The symbol | is read as "such that"

An **element** or **member** of a set is an object that belongs to the set

The symbol \in means "is an element of"

The symbol \notin means "is not an element of"

Generally capital letters are used to represent sets and lowercase letters are used for other objects i.e. $S=\{2,3,5,7\}$

Thus, $a \in S$ means a is an element of S

```
ls 2 \in \{0, 2, 4, 6\}?
ls 2 \in \{1, 3, 5, 7, 9\}?
```

An **element** or **member** of a set is an object that belongs to the set

The symbol \in means "is an element of"

The symbol \notin means "is not an element of"

Generally capital letters are used to represent sets and lowercase letters are used for other objects i.e. $S=\{2,3,5,7\}$

Thus, $a \in S$ means a is an element of S

```
\label{eq:second} \begin{array}{ll} \mathsf{Is} \ 2 \in \{0,2,4,6\}? & \mathsf{Yes} \\ \\ \mathsf{Is} \ 2 \in \{1,3,5,7,9\}? \end{array}
```

An **element** or **member** of a set is an object that belongs to the set

The symbol \in means "is an element of"

The symbol \notin means "is not an element of"

Generally capital letters are used to represent sets and lowercase letters are used for other objects i.e. $S=\{2,3,5,7\}$

Thus, $a \in S$ means a is an element of S

```
\label{eq:second} \begin{array}{ll} \mathsf{Is} \ 2 \in \{0,2,4,6\}? & \mathsf{Yes} \\ \\ \mathsf{Is} \ 2 \in \{1,3,5,7,9\}? & \mathsf{No}, \ 2 \notin \{1,3,5,7,9\} \end{array}
```

Some Important Sets

- \mathbb{N} Natural or Counting numbers: {1, 2, 3, ...}
- \mathbb{W} Whole Numbers: {0, 1, 2, 3, ...}
- \mathbb{I} Integers: {..., -3, -2, -1, 0, 1, 2, 3, ...}
- \mathbb{Q} Rational numbers: $\{\frac{p}{q} \mid p, q \in \mathbb{I}, q \neq 0\}$
- ℜ Real Numbers: { x | x is a number that can be written as a decimal }
- Irrational numbers: { x | x is a real number and x cannot be written as a quotient of integers }.
 Examples are: π, √2, and ³√4
- \emptyset Empty Set: { }, the set that contains nothing
- U Universal Set: the set of all objects currently under discussion

Any **rational** number can be written as either a **terminating** *decimal* (*like* 0.5, 0.333, or 0.8578966) *or a* **repeating** *decimal* (*like* 0.333 *or* 123.392545)

The decimal representation of an **irrational** number **never terminates** and **never repeats**

The set $\{ \emptyset \}$ is *not* empty, but is a set which *contains* the empty set

• Is
$$\emptyset \in \{ a, b, c \}$$
?

• Is
$$\frac{1}{3} \notin \{ x \mid x = \frac{1}{p}, p \in \mathbb{N} \}$$

More Membership Questions

• Is
$$\emptyset \in \{ a, b, c \}$$
? No

• Is
$$\emptyset \in \{ \emptyset, \{ \emptyset \} \}$$
?

• Is
$$\frac{1}{3} \notin \{ x \mid x = \frac{1}{p}, p \in \mathbb{N} \}$$

More Membership Questions

• Is
$$\emptyset \in \{ \mathsf{a}, \mathsf{b}, \mathsf{c} \}$$
? No

• Is
$$\frac{1}{3} \notin \{ x \mid x = \frac{1}{p}, p \in \mathbb{N} \}$$

More Membership Questions

- Is $\emptyset \in \{ a, b, c \}$? No
- Is $\emptyset \in \{ \emptyset, \{ \emptyset \} \}$? Yes
- Is $\emptyset \in \{ \{ \emptyset \} \}$? No
- Is $\frac{1}{3} \notin \{ x \mid x = \frac{1}{p}, p \in \mathbb{N} \}$

- Is $\emptyset \in \{ a, b, c \}$? No
- Is $\emptyset \in \{ \emptyset, \{ \emptyset \} \}$? Yes
- Is $\emptyset \in \{ \{ \emptyset \} \}$? No
- Is $\frac{1}{3} \notin \{ x \mid x = \frac{1}{p}, p \in \mathbb{N} \}$ No

The **cardinality** of a set is the number of **distinct** elements in the set

- $\bullet\,$ The cardinality of a set A is denoted ${\rm n}(A)$ or |A|
- If the cardinality of a set is a particular whole number, we call that set a **finite** set
- If a set is so large that there is no such number, it is called an **infinite** set (there is a precise definition of infinity but that is beyond the scope of this course)

Note: Sets do not care about the order or how many times an object is included. Thus, $\{1, 2, 3, 4\}$, $\{2, 3, 1, 4\}$, and $\{1, 2, 2, 3, 3, 3, 4, 4\}$ all describe the same set.

$$A = \{3, 5, 7, 9, 11\}, B = \{2, 4, 6, \dots, 100\}, C = \{1, 3, 5, 7, \dots\}$$
$$D = \{1, 2, 3, 2, 1\}, E = \{x \mid x \text{ is odd, and } x < 12\}$$

$$n(A) = ?$$

 $n(B) = ?$
 $n(C) = ?$
 $n(D) = ?$
 $n(E) = ?$

$$A = \{3, 5, 7, 9, 11\}, B = \{2, 4, 6, \dots, 100\}, C = \{1, 3, 5, 7, \dots\}$$
$$D = \{1, 2, 3, 2, 1\}, E = \{x \mid x \text{ is odd, and } x < 12\}$$

$$n(A) = ?$$
 5
 $n(B) = ?$
 $n(C) = ?$
 $n(D) = ?$
 $n(E) = ?$

$$A = \{3, 5, 7, 9, 11\}, B = \{2, 4, 6, \dots, 100\}, C = \{1, 3, 5, 7, \dots\}$$
$$D = \{1, 2, 3, 2, 1\}, E = \{x \mid x \text{ is odd, and } x < 12\}$$

$$\begin{array}{ll} {\sf n}(A) = ? & 5 \\ {\sf n}(B) = ? & 50 \\ {\sf n}(C) = ? & \\ {\sf n}(D) = ? & \\ {\sf n}(E) = ? & \end{array}$$

$$A = \{3, 5, 7, 9, 11\}, B = \{2, 4, 6, \dots, 100\}, C = \{1, 3, 5, 7, \dots\}$$
$$D = \{1, 2, 3, 2, 1\}, E = \{x \mid x \text{ is odd, and } x < 12\}$$

 $\begin{array}{ll} {\sf n}(A) = ? & 5 \\ {\sf n}(B) = ? & 50 \\ {\sf n}(C) = ? & {\sf Infinite} \\ {\sf n}(D) = ? \\ {\sf n}(E) = ? \end{array}$

$$A = \{3, 5, 7, 9, 11\}, B = \{2, 4, 6, \dots, 100\}, C = \{1, 3, 5, 7, \dots\}$$
$$D = \{1, 2, 3, 2, 1\}, E = \{x \mid x \text{ is odd, and } x < 12\}$$

 $\begin{array}{ll} {\sf n}(A) = ? & 5 \\ {\sf n}(B) = ? & 50 \\ {\sf n}(C) = ? & {\sf Infinite} \\ {\sf n}(D) = ? & 3 \\ {\sf n}(E) = ? \end{array}$

$$A = \{3, 5, 7, 9, 11\}, B = \{2, 4, 6, \dots, 100\}, C = \{1, 3, 5, 7, \dots\}$$
$$D = \{1, 2, 3, 2, 1\}, E = \{x \mid x \text{ is odd, and } x < 12\}$$

 $\begin{array}{ll} {\sf n}(A)=? & 5\\ {\sf n}(B)=? & 50\\ {\sf n}(C)=? & {\sf Infinite}\\ {\sf n}(D)=? & 3\\ {\sf n}(E)=? & 6 \end{array}$

- 1. every element of A is an element of B, and
- 2. every element of B is an element of A

$$\{ a, b, c \} = \{ b, c, a \} = \{ a, b, a, b, c \} ? \{ 3 \} = \{ x \mid x \in \mathbb{N} \text{ and } 1 < x < 5 \} ? \{ x \mid x \in \mathbb{N} \text{ and } x < 0 \} = \{ y \mid y \in \mathbb{Q} \text{ and } y \text{ is irrational} \} ?$$

- 1. every element of A is an element of B, and
- 2. every element of B is an element of A

$$\{ a, b, c \} = \{ b, c, a \} = \{ a, b, a, b, c \} ? Yes \{ 3 \} = \{ x \mid x \in \mathbb{N} \text{ and } 1 < x < 5 \} ? \{ x \mid x \in \mathbb{N} \text{ and } x < 0 \} = \{ y \mid y \in \mathbb{Q} \text{ and } y \text{ is irrational} \} ?$$

- 1. every element of A is an element of B, and
- 2. every element of B is an element of A

$$\{ a, b, c \} = \{ b, c, a \} = \{ a, b, a, b, c \} ? Yes \{ 3 \} = \{ x \mid x \in \mathbb{N} \text{ and } 1 < x < 5 \} ? No \{ x \mid x \in \mathbb{N} \text{ and } x < 0 \} = \{ y \mid y \in \mathbb{Q} \text{ and } y \text{ is irrational} \} ?$$

- 1. every element of A is an element of B, and
- 2. every element of B is an element of A

$$\{ a, b, c \} = \{ b, c, a \} = \{ a, b, a, b, c \} ? Yes \{ 3 \} = \{ x \mid x \in \mathbb{N} \text{ and } 1 < x < 5 \} ? No \{ x \mid x \in \mathbb{N} \text{ and } x < 0 \} = \{ y \mid y \in \mathbb{Q} \text{ and } y \text{ is irrational} \} ? Yes$$

Venn Diagrams & Subsets

Universe of Discourse – the set containing all elements under discussion for a particular problem

In mathematics, this is called the $\ensuremath{\mathbf{universal}}$ set and is denoted by U

Venn diagrams can be used to represent sets and their relationships to each other



The "Universe" is represented by the rectangle

Sets are represented with circles, shaded regions, and other shapes within the rectangle.



The Complement of a Set



The set A', the shaded region, is the complement of AA' is the set of all objects in the universe of discourse that are not elements of A

$$A' = \{x \mid x \in U \text{ and } x \notin A\}$$

Let
$$U = \{1, 2, 3, \dots, 8\}$$
, $R = \{1, 2, 5, 6\}$, and $S = \{2, 4, 5, 7, 8\}$

What is: R', the complement of R ?

What is: S', the complement of S ?

What is: U', the complement of U ?

What is: \emptyset' , the complement of \emptyset ?

Let
$$U = \{1, 2, 3, \dots, 8\}$$
, $R = \{1, 2, 5, 6\}$, and $S = \{2, 4, 5, 7, 8\}$

What is: R', the complement of R? $R' = \{3, 4, 7, 8\}$

What is: S', the complement of S ?

What is: U', the complement of U ?

What is: \emptyset' , the complement of \emptyset ?

Let
$$U = \{1, 2, 3, \dots, 8\}$$
, $R = \{1, 2, 5, 6\}$, and $S = \{2, 4, 5, 7, 8\}$

What is: R', the complement of R? $R' = \{3, 4, 7, 8\}$ What is: S', the complement of S? $S' = \{1, 3, 6\}$

What is: U', the complement of U ?

What is: \emptyset' , the complement of \emptyset ?

Let
$$U = \{1, 2, 3, \dots, 8\}$$
, $R = \{1, 2, 5, 6\}$, and $S = \{2, 4, 5, 7, 8\}$

What is: R', the complement of R? $R' = \{3, 4, 7, 8\}$ What is: S', the complement of S? $S' = \{1, 3, 6\}$ What is: U', the complement of U? $U' = \{\} = \emptyset$ What is: \emptyset' , the complement of \emptyset ?

Let
$$U = \{1, 2, 3, \dots, 8\}$$
, $R = \{1, 2, 5, 6\}$, and $S = \{2, 4, 5, 7, 8\}$

What is: R', the complement of R? $R' = \{3, 4, 7, 8\}$ What is: S', the complement of S? $S' = \{1, 3, 6\}$ What is: U', the complement of U? $U' = \{\} = \emptyset$ What is: \emptyset' , the complement of \emptyset ? $\emptyset' = U$

Set A is a **subset** of set B if every element of A is also an element of B. In other words, B contains all of the elements of A. This is denoted $A \subseteq B$.

Of the sets R, S, and T shown in the Venn diagram below, which are subsets?



Set A is a **subset** of set B if every element of A is also an element of B. In other words, B contains all of the elements of A. This is denoted $A \subseteq B$. Of the sets B. S and T shown in the Venn diagram below, which

Of the sets R, S, and T shown in the Venn diagram below, which are subsets?



 $T \subseteq R \text{ since } T \text{ is completely contained inside of } R$ Also every set is a subset of U

Suppose $U = \{1, 2, 3, \dots, 8\}$, $R = \{1, 2, 5, 6\}$, $S = \{2, 4, 5, 7, 8\}$, and $T = \{2, 6\}$

What element(s) are in the area where all the sets overlap?

What element(s) are in the area outside all the sets?



Suppose $U = \{1, 2, 3, \dots, 8\}$, $R = \{1, 2, 5, 6\}$, $S = \{2, 4, 5, 7, 8\}$, and $T = \{2, 6\}$

What element(s) are in the area where all the sets overlap?

What element(s) are in the area outside all the sets?



Another definition for set equality: Sets A and B are equal if and only if:

- 1. $A \subseteq B$ and
- **2**. $B \subseteq A$

Proper Subset: $A \subset B$ if $A \subseteq B$ and $A \neq B$
| $\{a,b,c\}$ | $\{a,c,d,f\}$ |
|-------------|-----------------|
| $\{a,b,c\}$ | $\{c,a,b\}$ |
| $\{a,b,c\}$ | $\{a,b,c\}$ |
| $\{a\}$ | $\{a,b,c\}$ |
| $\{a,c\}$ | $\{a,b,c,d\}$ |
| $\{a,c\}$ | $\{a,b,d,e,f\}$ |
| X | X |
| Ø | $\{a,b,c\}$ |
| Ø | Ø |

| $\{a, b, c\}$ | ¥ | $\{a,c,d,f\}$ |
|---------------|---|-----------------|
| $\{a,b,c\}$ | | $\{c,a,b\}$ |
| $\{a,b,c\}$ | | $\{a,b,c\}$ |
| $\{a\}$ | | $\{a,b,c\}$ |
| $\{a,c\}$ | | $\{a,b,c,d\}$ |
| $\{a,c\}$ | | $\{a,b,d,e,f\}$ |
| X | | X |
| Ø | | $\{a,b,c\}$ |
| Ø | | Ø |

| $\{a, b, c\}$ | ¥ | $\{a,c,d,f\}$ |
|---------------|-------------|-----------------|
| $\{a, b, c\}$ | \subseteq | $\{c,a,b\}$ |
| $\{a, b, c\}$ | | $\{a,b,c\}$ |
| $\{a\}$ | | $\{a,b,c\}$ |
| $\{a,c\}$ | | $\{a,b,c,d\}$ |
| $\{a,c\}$ | | $\{a,b,d,e,f\}$ |
| X | | X |
| Ø | | $\{a, b, c\}$ |
| Ø | | Ø |

| $\{a, b, c\}$ | ¥ | $\{a,c,d,f\}$ |
|---------------|-------------|-----------------|
| $\{a, b, c\}$ | \subseteq | $\{c,a,b\}$ |
| $\{a, b, c\}$ | \subseteq | $\{a,b,c\}$ |
| $\{a\}$ | | $\{a,b,c\}$ |
| $\{a,c\}$ | | $\{a,b,c,d\}$ |
| $\{a,c\}$ | | $\{a,b,d,e,f\}$ |
| X | | X |
| Ø | | $\{a,b,c\}$ |
| Ø | | Ø |

| $\{a, b, c\}$ | É | $\{a,c,d,f\}$ |
|---------------|-------------|-----------------|
| $\{a, b, c\}$ | \subseteq | $\{c,a,b\}$ |
| $\{a, b, c\}$ | \subseteq | $\{a,b,c\}$ |
| $\{a\}$ | \subset | $\{a,b,c\}$ |
| $\{a,c\}$ | | $\{a,b,c,d\}$ |
| $\{a,c\}$ | | $\{a,b,d,e,f\}$ |
| X | | X |
| Ø | | $\{a,b,c\}$ |
| Ø | | Ø |

| $\{a, b, c\}$ | É | $\{a,c,d,f\}$ |
|---------------|-------------|-----------------|
| $\{a, b, c\}$ | \subseteq | $\{c,a,b\}$ |
| $\{a, b, c\}$ | \subseteq | $\{a,b,c\}$ |
| $\{a\}$ | \subset | $\{a,b,c\}$ |
| $\{a,c\}$ | \subset | $\{a,b,c,d\}$ |
| $\{a,c\}$ | | $\{a,b,d,e,f\}$ |
| X | | X |
| Ø | | $\{a, b, c\}$ |
| Ø | | Ø |

Is the left set a subset of the set on the right?

 $\{a, b, c\} \not\subseteq \{a, c, d, f\}$ $\{a, b, c\} \subseteq \{c, a, b\}$ $\{a, b, c\} \subseteq \{a, b, c\}$ $\{a\} \subset \{a, b, c\}$ $\{a,c\} \subset \{a,b,c,d\}$ $\{a,c\} \not\subseteq \{a,b,d,e,f\}$ X = X $\emptyset \qquad \{a, b, c\}$ Ø Ø

Is the left set a subset of the set on the right?

 $\{a, b, c\} \not\subseteq \{a, c, d, f\}$ $\{a, b, c\} \subseteq \{c, a, b\}$ $\{a, b, c\} \subseteq \{a, b, c\}$ $\{a\} \subset \{a, b, c\}$ $\{a,c\} \subset \{a,b,c,d\}$ $\{a,c\} \not\subseteq \{a,b,d,e,f\}$ $X \subset X$ $\emptyset \qquad \{a, b, c\}$ Ø Ø

Is the left set a subset of the set on the right?

 $\{a, b, c\} \not\subseteq \{a, c, d, f\}$ $\{a, b, c\} \subseteq \{c, a, b\}$ $\{a, b, c\} \subseteq \{a, b, c\}$ $\{a\} \subset \{a, b, c\}$ $\{a,c\} \subset \{a,b,c,d\}$ $\{a,c\} \not\subseteq \{a,b,d,e,f\}$ $X \subset X$ $\emptyset \subset \{a, b, c\}$ Ø Ø

Is the left set a subset of the set on the right?

 $\{a, b, c\} \not\subseteq \{a, c, d, f\}$ $\{a, b, c\} \subseteq \{c, a, b\}$ $\{a, b, c\} \subseteq \{a, b, c\}$ $\{a\} \subset \{a, b, c\}$ $\{a,c\} \subset \{a,b,c,d\}$ $\{a,c\} \not\subseteq \{a,b,d,e,f\}$ $X \subset X$ $\emptyset \subset \{a, b, c\}$ $\emptyset \subseteq \emptyset$

- Any set is a subset of itself
- Any set is a subset of the universal set
- The empty set is a subset of every set including itself

| $\{1, 2, 3\}$ | I |
|-----------------|---------------------|
| $\{a,b\}$ | $\{a\}$ |
| $\{a\}$ | $\{a,b\}$ |
| $\{a,b,c\}$ | $\{a,d,e,g\}$ |
| $\{a,b,c\}$ | $\{a, a, c, b, c\}$ |
| $\{\emptyset\}$ | $\{a,b,c\}$ |
| $\{\emptyset\}$ | {} |

| $\{1, 2, 3\}$ | \subset | \mathbb{I} |
|-----------------|-----------|---------------------|
| $\{a,b\}$ | | $\{a\}$ |
| $\{a\}$ | | $\{a,b\}$ |
| $\{a,b,c\}$ | | $\{a,d,e,g\}$ |
| $\{a,b,c\}$ | | $\{a, a, c, b, c\}$ |
| $\{\emptyset\}$ | | $\{a,b,c\}$ |
| $\{\emptyset\}$ | | {} |

| $\{1, 2, 3\}$ | \subset | I |
|-----------------|-----------|---------------------|
| $\{a,b\}$ | ¢ | $\{a\}$ |
| $\{a\}$ | | $\{a,b\}$ |
| $\{a, b, c\}$ | | $\{a,d,e,g\}$ |
| $\{a,b,c\}$ | | $\{a, a, c, b, c\}$ |
| $\{\emptyset\}$ | | $\{a,b,c\}$ |
| $\{\emptyset\}$ | | {} |

| $\{1, 2, 3\}$ | \subset | I |
|-----------------|-----------|---------------------|
| $\{a,b\}$ | ¢ | $\{a\}$ |
| $\{a\}$ | \subset | $\{a,b\}$ |
| $\{a, b, c\}$ | | $\{a,d,e,g\}$ |
| $\{a,b,c\}$ | | $\{a, a, c, b, c\}$ |
| $\{\emptyset\}$ | | $\{a,b,c\}$ |
| $\{\emptyset\}$ | | {} |

 $\begin{array}{rcl} \{1,2,3\} &\subset & \mathbb{I} \\ & \{a,b\} & \not\subseteq & \{a\} \\ & \{a\} &\subset & \{a,b\} \\ \{a,b,c\} & \not\subseteq & \{a,d,e,g\} \\ & \{a,b,c\} & & \{a,a,c,b,c\} \\ & \{\emptyset\} & & \{a,b,c\} \\ & \{\emptyset\} & & \{\} \end{array}$

$$\begin{array}{rcl} \{1,2,3\} &\subset & \mathbb{I} \\ & \{a,b\} & \not\subseteq & \{a\} \\ & \{a\} &\subset & \{a,b\} \\ \{a,b,c\} & \not\subseteq & \{a,d,e,g\} \\ \{a,b,c\} &= & \{a,a,c,b,c\} \\ & \{\emptyset\} & & \{a,b,c\} \\ & \{\emptyset\} & & \{\} \end{array}$$

 $\begin{array}{rcl} \{1,2,3\} & \subset & \mathbb{I} \\ & \{a,b\} & \not\subseteq & \{a\} \\ & \{a\} & \subset & \{a,b\} \\ \{a,b,c\} & \not\subseteq & \{a,d,e,g\} \\ & \{a,b,c\} & = & \{a,a,c,b,c\} \\ & \{\emptyset\} & \not\subseteq & \{a,b,c\} \\ & \{\emptyset\} & \{\} \end{array}$

 $\begin{array}{rcl} \{1,2,3\} & \subset & \mathbb{I} \\ & \{a,b\} & \not\subseteq & \{a\} \\ & \{a\} & \subset & \{a,b\} \\ \{a,b,c\} & \not\subseteq & \{a,d,e,g\} \\ & \{a,b,c\} & = & \{a,a,c,b,c\} \\ & \{\emptyset\} & \not\subseteq & \{a,b,c\} \\ & \{\emptyset\} & \not\subseteq & \{\} \end{array}$

Find the following power sets and determine their cardinality.

$$\mathscr{P}(\emptyset) =$$

 $\mathscr{P}(\{a\}) =$
 $\mathscr{P}(\{a,b\}) =$
 $\mathscr{P}(\{a,b,c\}) =$

Find the following power sets and determine their cardinality.

| $\mathscr{P}(\emptyset) =$ | $\{\emptyset\}$ |
|----------------------------|-----------------|
| $\mathscr{P}(\{a\}) =$ | |
| $\mathscr{P}(\{a,b\}) =$ | |
| $\mathscr{P}(\{a,b,c\}) =$ | |

Find the following power sets and determine their cardinality.

| $\mathscr{P}(\emptyset) =$ | $\{\emptyset\}$ |
|----------------------------|--------------------------|
| $\mathscr{P}(\{a\}) =$ | $\{ \emptyset, \{a\} \}$ |
| $\mathscr{P}(\{a,b\}) =$ | |
| $\mathscr{P}(\{a,b,c\}) =$ | |

Find the following power sets and determine their cardinality.

| $\mathscr{P}(\emptyset) =$ | $\{\emptyset\}$ |
|----------------------------|-----------------|
|----------------------------|-----------------|

$$\mathscr{P}(\{a\}) = \quad \{ \emptyset, \{a\} \}$$

 $\mathscr{P}(\{a,b\}) = -\{\emptyset,\{a\},\{b\},\{a,b\}\}$

 $\mathscr{P}(\{a,b,c\}) =$

Find the following power sets and determine their cardinality.

$$\mathscr{P}(\emptyset) = \{\emptyset\}$$

$$\mathscr{P}(\{a\}) = \quad \{ \emptyset, \{a\} \}$$

 $\mathscr{P}(\{a,b\}) = -\{ \emptyset, \{a\}, \{b\}, \{a,b\} \}$

 $\mathscr{P}(\{a,b,c\}) = \quad \{ \emptyset, \{a\}, \{b\}, \{c\}, \{a,b\}, \{a,c\}, \{b,c\}, \{a,b,c\} \}$

Find the following power sets and determine their cardinality.

$$\mathscr{P}(\emptyset) = \{\emptyset\}$$

$$\mathscr{P}(\{a\}) = -\{ \emptyset, \{a\} \}$$

 $\mathscr{P}(\{a,b\}) = -\{ \emptyset, \{a\}, \{b\}, \{a,b\} \}$

 $\mathscr{P}(\{a,b,c\}) = \quad \{ \emptyset, \{a\}, \{b\}, \{c\}, \{a,b\}, \{a,c\}, \{b,c\}, \{a,b,c\} \}$

Is there a pattern?

 $1, 2, 4, 8, 16, 32, 64, \ldots, 2^n, \ldots$

Another Method for Generating Power Sets

A tree diagram can be used to generate $\mathscr{P}(A)$. Each element of the set is either in a particular subset, or it's not.



The number of subsets of a set with cardinality n is 2^n The number of **proper** subsets is $2^n - 1$ (Why?)

Intersection

The intersection of two sets, $A \cap B$, is the set of elements common to both: $A \cap B = \{x | x \in A \text{ and } x \in B\}$.

In other words, for an object to be in $A \cap B$ it must be a member of both A and B.



Intersection

The intersection of two sets, $A \cap B$, is the set of elements common to both: $A \cap B = \{x | x \in A \text{ and } x \in B\}$.

In other words, for an object to be in $A \cap B$ it must be a member of both A and B.



 $\{a, b, c\} \cap \{b, f, g\} =$ $\{a, b, c\} \cap \{a, b, c\} =$ For any $A, A \cap A =$ $\{a, b, c\} \cap \{a, b, z\} =$ $\{a, b, c\} \cap \{x, y, z\} =$ $\{a, b, c\} \cap \emptyset =$ = For any $A, A \cap \emptyset$ = For any $A, A \cap U$ For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, g\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} =$ For any $A, A \cap A =$ $\{a, b, c\} \cap \{a, b, z\} =$ $\{a, b, c\} \cap \{x, y, z\} =$ $\{a, b, c\} \cap \emptyset =$ For any $A, A \cap \emptyset$ = For any $A, A \cap U$ = For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, g\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any A, $A \cap A =$ $\{a, b, c\} \cap \{a, b, z\} =$ $\{a, b, c\} \cap \{x, y, z\} =$ $\{a, b, c\} \cap \emptyset =$ For any $A, A \cap \emptyset$ = For any $A, A \cap U$ = For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, q\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} =$ $\{a, b, c\} \cap \{x, y, z\} =$ $\{a, b, c\} \cap \emptyset =$ For any $A, A \cap \emptyset$ = = For any $A, A \cap U$ For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, q\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} = \{a, b\}$ $\{a, b, c\} \cap \{x, y, z\} =$ $\{a, b, c\} \cap \emptyset =$ For any $A, A \cap \emptyset =$ For any $A, A \cap U$ = For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, g\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} = \{a, b\}$ $\{a, b, c\} \cap \{x, y, z\} = \emptyset$ $\{a, b, c\} \cap \emptyset =$ For any $A, A \cap \emptyset =$ For any $A, A \cap U$ = For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, q\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} = \{a, b\}$ $\{a, b, c\} \cap \{x, y, z\} = \emptyset$ $\{a, b, c\} \cap \emptyset = \emptyset$ For any $A, A \cap \emptyset =$ For any $A, A \cap U$ = For any $A \subseteq B$, $A \cap B =$

 $\{a, b, c\} \cap \{b, f, q\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} = \{a, b\}$ $\{a, b, c\} \cap \{x, y, z\} = \emptyset$ $\{a, b, c\} \cap \emptyset = \emptyset$ For any $A, A \cap \emptyset = \emptyset$ For any $A, A \cap U$ = For any $A \subseteq B$, $A \cap B =$
Find the Following Intersections

 $\{a, b, c\} \cap \{b, f, g\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} = \{a, b\}$ $\{a, b, c\} \cap \{x, y, z\} = \emptyset$ $\{a, b, c\} \cap \emptyset = \emptyset$ For any $A, A \cap \emptyset = \emptyset$ For any $A, A \cap U = A$ For any $A \subseteq B$, $A \cap B =$

Find the Following Intersections

 $\{a, b, c\} \cap \{b, f, g\} = \{b\}$ $\{a, b, c\} \cap \{a, b, c\} = \{a, b, c\}$ For any $A, A \cap A = A$ $\{a, b, c\} \cap \{a, b, z\} = \{a, b\}$ $\{a, b, c\} \cap \{x, y, z\} = \emptyset$ $\{a, b, c\} \cap \emptyset = \emptyset$ For any $A, A \cap \emptyset = \emptyset$ For any $A, A \cap U = A$ For any $A \subseteq B$, $A \cap B = A$

Disjoint sets: two sets which have no elements in common.

I.e., their intersection is empty: $A \cap B = \emptyset$



$$\left\{ \begin{array}{l} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{l} d, e, f, g \end{array} \right\} \\ \left\{ \begin{array}{l} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{l} a, b, c \end{array} \right\} \\ \left\{ \begin{array}{l} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{l} a, b, z \end{array} \right\} \\ \left\{ \begin{array}{l} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{l} x, y, z \end{array} \right\} \\ \left\{ \begin{array}{l} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{l} x, y, z \end{array} \right\} \\ \left\{ \begin{array}{l} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \end{array} \right\} \\ For any A, A and \emptyset \\ For any A, A and A' \end{array}$$

$$\{a, b, c\}$$
 and $\{d, e, f, g\}$ Yes

- $\{ \text{ a, b, c} \} \quad \text{and} \quad \{ \text{ a, b, c} \}$
- $\{ \text{ a, b, c} \} \quad \text{and} \quad \{ \text{ a, b, z} \}$
- $\left\{ \text{ a, b, c} \right\} \quad \text{and} \quad \left\{ \text{ x, y, z} \right\}$
- $\{a, b, c\}$ and \emptyset

For any A, A and \emptyset

$$\{a, b, c\}$$
 and $\{d, e, f, g\}$ Yes

$$\{a, b, c\}$$
 and $\{a, b, c\}$ No

$$\{ \text{ a, b, c} \} \quad \text{and} \quad \{ \text{ a, b, z} \}$$

$$\{ \text{ a, b, c} \} \quad \text{and} \quad \{ \text{ x, y, z} \}$$

$$\{a, b, c\}$$
 and \emptyset

For any $A,\,A$ and \emptyset

$$\{ \ \mathsf{a}, \ \mathsf{b}, \ \mathsf{c} \ \} \quad \mathsf{and} \quad \{ \ \mathsf{d}, \ \mathsf{e}, \ \mathsf{f}, \ \mathsf{g} \ \} \quad \begin{tabular}{c} \end{tabular} \mathbf{Yes} \\ \end{tabular}$$

$$\{a, b, c\}$$
 and $\{a, b, c\}$ No

- $\{a, b, c\}$ and $\{a, b, z\}$ No
- $\left\{ \text{ a, b, c} \right\} \quad \text{and} \quad \left\{ \text{ x, y, z} \right\}$
- $\{ \text{ a, b, c} \} \quad \text{and} \quad \emptyset$

For any A, A and \emptyset

$${a, b, c}$$
 and ${d, e, f, g}$ Yes
 ${a, b, c}$ and ${a, b, c}$ No

 $\left\{ \text{ a, b, c} \right\} \quad \text{and} \quad \left\{ \text{ a, b, z} \right\} \qquad \textbf{No}$

$$\{ a, b, c \}$$
 and $\{ x, y, z \}$ Yes

$$\{a, b, c\}$$
 and \emptyset

For any A, A and \emptyset

$$\left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} d, e, f, g \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{No} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} a, b, z \end{array} \right\} \quad \text{No} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} a, b, c \end{array} \right\} \quad \text{and} \quad \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}{c} x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, y, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x] x, z \end{array} \right\} \quad \text{Yes} \\ \left\{ \begin{array}[x]$$

| $\{ a, b, c \}$ | and | $\left\{ \text{ d, e, f, g} \right\}$ | Yes | | |
|-----------------------------------|-----|---------------------------------------|-----|--|--|
| $\{ a, b, c \}$ | and | $\{ a, b, c \}$ | Νο | | |
| $\{ a, b, c \}$ | and | $\{ a, b, z \}$ | Νο | | |
| $\{a, b, c\}$ | and | $\{ x, y, z \}$ | Yes | | |
| $\{a, b, c\}$ | and | Ø | Yes | | |
| For any A , A and \emptyset | | | Yes | | |
| For any A , A and A' | | | | | |

| $\{ a, b, c \}$ | and | $\left\{ \text{ d, e, f, g} \right\}$ | Yes |
|---------------------------------------------------------|-----|---------------------------------------|-----|
| $\{ a, b, c \}$ | and | $\{ a, b, c \}$ | Νο |
| $\{a, b, c\}$ | and | $\{ \text{ a, b, z } \}$ | Νο |
| $\{a, b, c\}$ | and | $\{ x, y, z \}$ | Yes |
| $\{ a, b, c \}$ | and | Ø | Yes |
| For any A , | Yes | | |
| For any A,A and \emptyset For any A,A and A' | | Yes | |

Set Union

The union of two sets, $A \cup B$, is the set of elements belonging to either of the sets: $A \cup B = \{x | x \in A \text{ or } x \in B\}$

In other words, for an object to be in $A \cup B$ it must be a member of either A or B.



Set Union

The union of two sets, $A \cup B$, is the set of elements belonging to either of the sets: $A \cup B = \{x | x \in A \text{ or } x \in B\}$

In other words, for an object to be in $A \cup B$ it must be a member of either A or B.



$$\{ \text{ a, b, c} \} \hspace{0.1in} \cup \hspace{0.1in} \{ \hspace{0.1in} \text{b, f, g} \hspace{0.1in} \} \hspace{0.1in} = \hspace{0.1in}$$

$$\{ a, b, c \} \cup \{ a, b, c \} =$$

For any A, $A \cup A =$

$$\{ \text{ a, b, c} \} \hspace{0.1in} \cup \hspace{0.1in} \{ \text{ a, b, z} \} \hspace{0.1in} = \hspace{0.1in}$$

$$\{ \text{ a, b, c} \} \ \cup \ \{ \text{ x, y, z} \} \ =$$

$$\{ \text{ a, b, c} \} \hspace{0.1 in} \cup \hspace{0.1 in} \emptyset \hspace{0.1 in} = \hspace{0.1 in}$$

For any $A, A \cup \emptyset =$

For any $A, A \cup U =$

For any $A \subseteq B$, $A \cup B =$

$$\left\{ \begin{array}{l} \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \begin{array}{l} \mathsf{b}, \mathsf{f}, \mathsf{g} \right\} &= \left\{ a, b, c, f, g \right\} \\ \left\{ \begin{array}{l} \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} &= \\ \end{array} \right. \\ \left\{ \begin{array}{l} \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} &= \\ \left\{ \begin{array}{l} \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{a}, \mathsf{b}, \mathsf{z} \right\} &= \\ \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{x}, \mathsf{y}, \mathsf{z} \right\} &= \\ \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left\{ \begin{array}{l} \mathsf{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \\ \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \\ \left. \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left. \left\{ \operatorname{a}, \mathsf{b}, \mathsf{c} \right\} \cup \left\{ \mathsf{w} \right\} &= \\ \end{array} \\ \left. \left\{ \operatorname{a}, \mathsf{w}, \mathsf{w} \right\} &= \\ \left. \left\{ \operatorname{a}, \mathsf{w}, \mathsf{w} \right\} &= \\ \end{array} \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \end{array} \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \end{array} \right. \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &= \\ \left. \left\{ \operatorname{a}, \mathsf{w} \right\} &=$$

For any $A \subseteq B$, $A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A =$

$$\{a, b, c\} \cup \{a, b, z\} =$$

$$\{a, b, c\} \cup \{x, y, z\} =$$

$$\{a, b, c\} \cup \emptyset =$$
For any $A, A \cup \emptyset =$
For any $A, A \cup U =$
For any $A, A \cup U =$
For any $A, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$
$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$
$$\{a, b, c\} \cup \{a, b, z\} =$$
$$\{a, b, c\} \cup \{x, y, z\} =$$
$$\{a, b, c\} \cup \emptyset =$$
For any $A, A \cup \emptyset =$ For any $A, A \cup U =$ For any $A, A \cup U =$ For any $A \subseteq B, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$

$$\{a, b, c\} \cup \{a, b, z\} = \{a, b, c, z\}$$

$$\{a, b, c\} \cup \{x, y, z\} =$$

$$\{a, b, c\} \cup \emptyset =$$
For any $A, A \cup \emptyset =$
For any $A, A \cup U =$
For any $A, A \cup U =$
For any $A, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$

$$\{a, b, c\} \cup \{a, b, z\} = \{a, b, c, z\}$$

$$\{a, b, c\} \cup \{x, y, z\} = \{a, b, c, x, y, z\}$$

$$\{a, b, c\} \cup \emptyset =$$
For any $A, A \cup \emptyset =$
For any $A, A \cup U =$
For any $A, A \cup U =$
For any $A \subseteq B, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$

$$\{a, b, c\} \cup \{a, b, z\} = \{a, b, c, z\}$$

$$\{a, b, c\} \cup \{x, y, z\} = \{a, b, c, x, y, z\}$$

$$\{a, b, c\} \cup \emptyset = \{a, b, c\}$$
For any $A, A \cup \emptyset =$
For any $A, A \cup U =$
For any $A, A \cup U =$
For any $A \subseteq B, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$

$$\{a, b, c\} \cup \{a, b, z\} = \{a, b, c, z\}$$

$$\{a, b, c\} \cup \{x, y, z\} = \{a, b, c, x, y, z\}$$

$$\{a, b, c\} \cup \emptyset = \{a, b, c\}$$
For any $A, A \cup \emptyset = A$
For any $A, A \cup U =$
For any $A, A \cup U =$
For any $A \subseteq B, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$

$$\{a, b, c\} \cup \{a, b, z\} = \{a, b, c, z\}$$

$$\{a, b, c\} \cup \{x, y, z\} = \{a, b, c, x, y, z\}$$

$$\{a, b, c\} \cup \emptyset = \{a, b, c\}$$
For any $A, A \cup \emptyset = A$
For any $A, A \cup U = U$
For any $A, A \cup U = U$
For any $A \subseteq B, A \cup B =$

$$\{a, b, c\} \cup \{b, f, g\} = \{a, b, c, f, g\}$$

$$\{a, b, c\} \cup \{a, b, c\} = \{a, b, c\}$$
For any $A, A \cup A = A$

$$\{a, b, c\} \cup \{a, b, z\} = \{a, b, c, z\}$$

$$\{a, b, c\} \cup \{x, y, z\} = \{a, b, c, x, y, z\}$$

$$\{a, b, c\} \cup \emptyset = \{a, b, c\}$$
For any $A, A \cup \emptyset = A$
For any $A, A \cup U = U$
For any $A \subseteq B, A \cup B = B$

Set Difference

The difference of two sets, A - B, is the set of elements belonging to set A and not to set B: $A - B = \{x | x \in A \text{ and } x \notin B\}$



Note: $x \notin B$ means $x \in B'$ Thus, $A - B = \{x | x \in A \text{ and } x \in B'\}$ $= A \cap B'$

Set Difference

The difference of two sets, A - B, is the set of elements belonging to set A and not to set B: $A - B = \{x | x \in A \text{ and } x \notin B\}$



Note: $x \notin B$ means $x \in B'$ Thus, $A - B = \{x | x \in A \text{ and } x \in B'\}$ $= A \cap B'$

$$\{1, 2, 3, 4, 5\} - \{2, 4, 6\} =$$

$$\{2,4,6\} - \{1,2,3,4,5\} =$$

Note, in general, $A - B \neq B - A$

$$\{1, 2, 3, 4, 5\} - \{2, 4, 6\} = \{1, 3, 5\}$$

$$\{2,4,6\} - \{1,2,3,4,5\} =$$

Note, in general, $A - B \neq B - A$

$$\{1, 2, 3, 4, 5\} - \{2, 4, 6\} = \{1, 3, 5\}$$

$$\{2,4,6\} - \{1,2,3,4,5\} = \{6\}$$

Note, in general, $A - B \neq B - A$

 $U = \{1, 2, 3, 4, 5, 6, 9\}$ $A = \{1, 2, 3, 4\}$ $B = \{2, 4, 6\}$ $C = \{1, 3, 6, 9\}$

Find each of these sets:

- $A \cup B =$
- $\bullet \ A \cap B =$
- $A \cap U =$

$\bullet \ A \cup U =$

 $U = \{1, 2, 3, 4, 5, 6, 9\}$ $A = \{1, 2, 3, 4\}$ $B = \{2, 4, 6\}$ $C = \{1, 3, 6, 9\}$

Find each of these sets:

- $A \cup B = \{1, 2, 3, 4, 6\}$
- $\bullet \ A \cap B =$
- $\bullet \ A \cap U =$

$\bullet \ A \cup U =$

 $U = \{1, 2, 3, 4, 5, 6, 9\}$ $A = \{1, 2, 3, 4\}$ $B = \{2, 4, 6\}$ $C = \{1, 3, 6, 9\}$

Find each of these sets:

- $A \cup B = \{1, 2, 3, 4, 6\}$
- $A \cap B = \{2, 4\}$
- $\bullet \ A \cap U =$

$\bullet \ A \cup U =$

 $U = \{1, 2, 3, 4, 5, 6, 9\}$ $A = \{1, 2, 3, 4\}$ $B = \{2, 4, 6\}$ $C = \{1, 3, 6, 9\}$

Find each of these sets:

- $A \cup B = \{1, 2, 3, 4, 6\}$
- $A \cap B = \{2, 4\}$
- $A \cap U = A = \{1, 2, 3, 4\}$

• $A \cup U =$

 $U = \{1, 2, 3, 4, 5, 6, 9\}$ $A = \{1, 2, 3, 4\}$ $B = \{2, 4, 6\}$ $C = \{1, 3, 6, 9\}$

- $A \cup B = \{1, 2, 3, 4, 6\}$
- $A \cap B = \{2, 4\}$
- $A \cap U = A = \{1, 2, 3, 4\}$
- $A \cup U = U = \{1, 2, 3, 4, 5, 6, 9\}$

$$U = \{1, 2, 3, 4, 5, 6, 9\}$$

$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 3, 6, 9\}$$

•
$$A' \cap B =$$

- $A' \cup B =$
- $A \cup B \cup C =$

•
$$A \cap B \cap C =$$

$$U = \{1, 2, 3, 4, 5, 6, 9\}$$

$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 3, 6, 9\}$$

•
$$A' = \{5, 6, 9\}$$

•
$$A' \cap B =$$

- $A' \cup B =$
- $A \cup B \cup C =$

•
$$A \cap B \cap C =$$

$$U = \{1, 2, 3, 4, 5, 6, 9\}$$

$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 3, 6, 9\}$$

•
$$A' = \{5, 6, 9\}$$

•
$$A' \cap B = \{6\}$$

- $A' \cup B =$
- $A \cup B \cup C =$

•
$$A \cap B \cap C =$$
$$U = \{1, 2, 3, 4, 5, 6, 9\}$$

$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 3, 6, 9\}$$

Find each of these sets:

•
$$A' = \{5, 6, 9\}$$

•
$$A' \cap B = \{6\}$$

•
$$A' \cup B = \{5, 6, 9, 2, 4\}$$

•
$$A \cup B \cup C =$$

•
$$A \cap B \cap C =$$

$$U = \{1, 2, 3, 4, 5, 6, 9\}$$

$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 3, 6, 9\}$$

Find each of these sets:

•
$$A' = \{5, 6, 9\}$$

•
$$A' \cap B = \{6\}$$

•
$$A' \cup B = \{5, 6, 9, 2, 4\}$$

•
$$A \cup B \cup C = \{1, 2, 3, 4, 6, 9\}$$

 $\bullet \ A \cap B \cap C =$

$$U = \{1, 2, 3, 4, 5, 6, 9\}$$

$$A = \{1, 2, 3, 4\}$$

$$B = \{2, 4, 6\}$$

$$C = \{1, 3, 6, 9\}$$

Find each of these sets:

•
$$A' = \{5, 6, 9\}$$

•
$$A' \cap B = \{6\}$$

•
$$A' \cup B = \{5, 6, 9, 2, 4\}$$

•
$$A \cup B \cup C = \{1, 2, 3, 4, 6, 9\}$$

• $A \cap B \cap C = \emptyset$

•
$$A' \cup B' =$$

•
$$A' \cap B' =$$

•
$$A \cap (B \cup C) =$$

•
$$(A' \cup C) \cap B =$$

• $A' \cup B' =$ Not in A or not in B

•
$$A' \cap B' =$$

•
$$A \cap (B \cup C) =$$

•
$$(A' \cup C) \cap B =$$

•
$$A' \cup B' =$$
Not in A or not in B

•
$$A' \cap B' =$$
 Not in A and not in B

•
$$A \cap (B \cup C) =$$

•
$$(A' \cup C) \cap B =$$

•
$$A' \cup B' =$$
Not in A or not in B

•
$$A' \cap B' =$$
 Not in A and not in B

• $A \cap (B \cup C) = \ln A$ and either in B or in C (or both)

•
$$(A' \cup C) \cap B =$$

•
$$A' \cup B' =$$
Not in A or not in B

•
$$A' \cap B' =$$
 Not in A and not in B

• $A \cap (B \cup C) = \ln A$ and either in B or in C (or both)

• $(A' \cup C) \cap B =$ In *B* and either not in *A* or in *C*

$$U = \{1, 2, 3, 4, 5, 6, 7\}$$

$$A = \{1, 2, 3, 4, 5, 6\}$$

$$B = \{2, 3, 6\}$$

$$C = \{3, 5, 7\}$$

Find each set:

• A - B =

•
$$B - A =$$

•
$$(A-B) \cup C' =$$

$$U = \{1, 2, 3, 4, 5, 6, 7\}$$

$$A = \{1, 2, 3, 4, 5, 6\}$$

$$B = \{2, 3, 6\}$$

$$C = \{3, 5, 7\}$$

Find each set:

- $A B = \{1, 4, 5\}$
- $\bullet \ B-A =$

•
$$(A-B) \cup C' =$$

$$U = \{1, 2, 3, 4, 5, 6, 7\}$$

$$A = \{1, 2, 3, 4, 5, 6\}$$

$$B = \{2, 3, 6\}$$

$$C = \{3, 5, 7\}$$

Find each set:

- $A B = \{1, 4, 5\}$
- $B A = \emptyset$

•
$$(A-B) \cup C' =$$

$$U = \{1, 2, 3, 4, 5, 6, 7\}$$

$$A = \{1, 2, 3, 4, 5, 6\}$$

$$B = \{2, 3, 6\}$$

$$C = \{3, 5, 7\}$$

Find each set:

• $A - B = \{1, 4, 5\}$

•
$$B - A = \emptyset$$

•
$$(A - B) \cup C' = \{1, 4, 5, 2, 6\}$$

Finding intersections, unions, differences, and complements of sets are examples of set operations

An **operation** is a rule or procedure by which one or more objects are used to obtain another object (usually a set or number).

Common Set Operations

Let A and B be any sets, with U the universal set.

• The complement of A is: $A' = \{x | x \in U \text{ and } x \notin A\}$



Set Intersection and Union

• The intersection of A and B is: $A \cap B = \{x | x \in A \text{ and } x \in B\}$



• The union of A and B is: $A \cup B = \{x | x \in A \text{ or } x \in B\}$



The difference of A and B is: $A - B = \{x | x \in A \text{ and } x \notin B\}$



Suppose
$$U = \{q, r, s, t, u, v, w, x, y, z\}$$
,
 $A = \{r, s, t, i, v\}$,
and $B = \{t, v, x\}$

Complete the Venn Diagram to represent $U,\,A,\,{\rm and}\,\,B$



Suppose
$$U = \{q, r, s, t, u, v, w, x, y, z\}$$
,
 $A = \{r, s, t, i, v\}$,
and $B = \{t, v, x\}$

Complete the Venn Diagram to represent U, A, and B



Shade the Diagram for: $A' \cap B' \cap C$



Shade the Diagram for: $A' \cap B' \cap C$



Shade the Diagram for: $(A \cap B)'$



Shade the Diagram for: $A' \cup B'$



Shade the Diagram for: $(A \cap B)'$



Shade the Diagram for: $A' \cup B'$



Shade the Diagram for: $(A \cap B)'$



Shade the Diagram for: $A' \cup B'$



We get the same region. $(A \cap B)' = A' \cup B'$

De Morgan's Laws

De Morgan's Laws: For any sets A and B

•
$$(A \cap B)' = A' \cup B'$$

• $(A \cup B)' = A' \cap B'$

Using A, B, C, \cap , \cup , -, and ', give a symbolic description of the shaded area in each of the following diagrams. Is there more than one way to describe each?





Cardinal Numbers & Surveys

Suppose,

 $\label{eq:constraint} \begin{array}{l} U = \mbox{The set of all students at EIU} \\ A = \mbox{The set of all male 2120 students} \\ B = \mbox{The set of all female 2120 students} \\ A \cup B = \end{array}$

Now suppose,

n(A) = 97n(B) = 101



$$n(A \cup B) =$$

Cardinal Numbers & Surveys

Suppose,

U = The set of all students at EIU A = The set of all male 2120 students B = The set of all female 2120 students $A \cup B =$ **The set of all 2120 students**

Now suppose,

n(A) = 97n(B) = 101



$$n(A \cup B) =$$

Cardinal Numbers & Surveys

Suppose,

U = The set of all students at EIU A = The set of all male 2120 students B = The set of all female 2120 students $A \cup B =$ **The set of all 2120 students**

Now suppose,

n(A) = 97n(B) = 101



$$n(A \cup B) = 97 + 101 = 198$$

U = The set of all students at EIU A = The set of all 2120 students that own a car B = The set of all 2120 students that own a truck $A \cup B =$ $A \cap B =$



U = The set of all students at EIU A = The set of all 2120 students that own a car B = The set of all 2120 students that own a truck $A \cup B =$ All 2120 students that own a car or a truck $A \cap B =$



U = The set of all students at EIU A = The set of all 2120 students that own a car B = The set of all 2120 students that own a truck $A \cup B =$ All 2120 students that own a car or a truck $A \cap B =$ All 2120 students that own a car and a truck



U = The set of all students at EIU A = The set of all 2120 students that own a car B = The set of all 2120 students that own a truck $A \cup B =$ All 2120 students that own a car or a truck $A \cap B =$ All 2120 students that own a car and a truck



U = The set of all students at EIU A = The set of all 2120 students that own a car B = The set of all 2120 students that own a truck $A \cup B =$ All 2120 students that own a car or a truck $A \cap B =$ All 2120 students that own a car and a truck

Now suppose, n(A) = 33 n(B) = 27 $n(A \cap B) = 10$ $n(A \cup B) = 23 + 10 + 17 = 50$



U = The set of all students at EIU A = The set of all 2120 students that own a car B = The set of all 2120 students that own a truck $A \cup B =$ All 2120 students that own a car or a truck $A \cap B =$ All 2120 students that own a car and a truck

Now suppose, n(A) = 33 n(B) = 27 $n(A \cap B) = 10$ $n(A \cup B) = 23 + 10 + 17 = 50$ = 33 + 27 - 10



For any two finite sets A and B: $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

In other words, the number of elements in the union of two sets is the sum of the number of elements in each of the sets minus the number of elements in their intersection.

How many integers between 1 and 100 are divisible by 2 or 5?

Let,

 $\begin{array}{lll} A=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }2\}\\ B=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }5\}\\ n(A)=&\\ n(B)=&\\ n(A\cap B)=&\\ n(A\cup B)=& \end{array}$

For any two finite sets A and B: $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

In other words, the number of elements in the union of two sets is the sum of the number of elements in each of the sets minus the number of elements in their intersection.

How many integers between 1 and 100 are divisible by 2 or 5?

Let,

 $\begin{array}{lll} A=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }2\}\\ B=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }5\}\\ n(A)=&\mathbf{50}\\ n(B)=&\\ n(A\cap B)=&\\ n(A\cup B)=& \end{array}$

For any two finite sets A and B: $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

In other words, the number of elements in the union of two sets is the sum of the number of elements in each of the sets minus the number of elements in their intersection.

How many integers between 1 and 100 are divisible by 2 or 5?

Let,

 $\begin{array}{rll} A=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }2\}\\ B=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }5\}\\ n(A)=&\textbf{50}\\ n(B)=&\textbf{20}\\ n(A\cap B)=&\\ n(A\cup B)=&\end{array}$

For any two finite sets A and B: $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

In other words, the number of elements in the union of two sets is the sum of the number of elements in each of the sets minus the number of elements in their intersection.

How many integers between 1 and 100 are divisible by 2 or 5?

Let,

 $\begin{array}{rcl} A=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }2\}\\ B=&\{n\mid 1\leq n\leq 100 \text{ and }n \text{ is divisible by }5\}\\ n(A)=&\mathbf{50}\\ n(B)=&\mathbf{20}\\ n(A\cap B)=&\mathbf{10}\\ n(A\cup B)=&\end{array}$
Inclusion/Exclusion Principle

For any two finite sets A and B: $n(A \cup B) = n(A) + n(B) - n(A \cap B)$

In other words, the number of elements in the union of two sets is the sum of the number of elements in each of the sets minus the number of elements in their intersection.

How many integers between 1 and 100 are divisible by 2 or 5?

Let,

 $\begin{array}{rll} A = & \{n \mid 1 \leq n \leq 100 \text{ and } n \text{ is divisible by } 2\} \\ B = & \{n \mid 1 \leq n \leq 100 \text{ and } n \text{ is divisible by } 5\} \\ n(A) = & \mathbf{50} \\ n(B) = & \mathbf{20} \\ n(A \cap B) = & \mathbf{10} \\ n(A \cup B) = & \mathbf{50} + \mathbf{20} - \mathbf{10} = \mathbf{60} \end{array}$



$$\begin{array}{ccc} A & (A \cup B) \cap C \\ B & A' \\ A \cap B \cap C' & C - B \\ A \cup B & (A \cup B) \cap C' \end{array}$$



$$A \qquad \mathbf{18} \quad (A \cup B) \cap C$$
$$B \qquad A'$$
$$A \cap B \cap C' \qquad C - B$$
$$A \cup B \qquad (A \cup B) \cap C'$$



$$A \qquad \mathbf{18} \quad (A \cup B) \cap C$$
$$B \qquad \mathbf{6} \qquad A'$$
$$A \cap B \cap C' \qquad C - B$$
$$A \cup B \qquad (A \cup B) \cap C'$$



$$A \qquad \mathbf{18} \qquad (A \cup B) \cap C$$
$$B \qquad \mathbf{6} \qquad A'$$
$$A \cap B \cap C' \qquad \mathbf{3} \qquad C - B$$
$$A \cup B \qquad (A \cup B) \cap C'$$



$$\begin{array}{cccc} A & 18 & (A \cup B) \cap C \\ B & 6 & A' \\ A \cap B \cap C' & 3 & C - B \\ A \cup B & 20 & (A \cup B) \cap C' \end{array}$$



$$\begin{array}{ccccc}
A & 18 & (A \cup B) \cap C & 5 \\
B & 6 & A' \\
A \cap B \cap C' & 3 & C - B \\
A \cup B & 20 & (A \cup B) \cap C' \end{array}$$



Find the cardinality of the sets:

 $\begin{array}{ccccc}
A & 18 & (A \cup B) \cap C & 5 \\
B & 6 & A' & 9 \\
A \cap B \cap C' & 3 & C - B \\
A \cup B & 20 & (A \cup B) \cap C' \\
\end{array}$



Find the cardinality of the sets:

- $A \qquad \mathbf{18} \qquad (A \cup B) \cap C \qquad \mathbf{5}$
- B 6 A' 9
- $A \cap B \cap C'$ **3** C B **6**

 $A \cup B$ **20** $(A \cup B) \cap C'$



- $A \qquad \mathbf{18} \quad (A \cup B) \cap C \qquad \mathbf{5}$
- B 6 A' 9
- $A \cap B \cap C'$ **3** C B **6**
 - $A \cup B$ **20** $(A \cup B) \cap C'$ **15**

Venn Diagram for 2 Sets

There are four disjoint regions



- I: $A \cap B'$
- II: $A \cap B$
- ${\sf III:} \quad A' \quad \cap \quad B$
- $\mathsf{IV:} \quad A' \quad \cap \quad B'$

There are eight disjoint regions



Technique for Counting with Venn Diagrams

- Designate the universal set
- Describe the sets of interest
- Draw a general Venn diagram
- Relate known information to the sizes of the disjoint regions of the diagram
- Infer the sizes of any remaining regions

- 5 on which both Simon and Garfunkel sing
- 8 total on which Simon sings
- 7 total on which Garfunkel sings
- 12 on which neither Simon nor Garfunkel sings



- 1. How many of her CDs feature only Paul Simon?
- 2. How many of her CDs feature only Art Garfunkel?
- 3. How many feature at least one of these two artists?

- 5 on which both Simon and Garfunkel sing
- 8 total on which Simon sings
- 7 total on which Garfunkel sings
- 12 on which neither Simon nor Garfunkel sings



- 1. How many of her CDs feature only Paul Simon?
- 2. How many of her CDs feature only Art Garfunkel?
- 3. How many feature at least one of these two artists?

- 5 on which both Simon and Garfunkel sing
- 8 total on which Simon sings
- 7 total on which Garfunkel sings
- 12 on which neither Simon nor Garfunkel sings



- 1. How many of her CDs feature only Paul Simon? 3
- 2. How many of her CDs feature only Art Garfunkel?
- 3. How many feature at least one of these two artists?

- 5 on which both Simon and Garfunkel sing
- 8 total on which Simon sings
- 7 total on which Garfunkel sings
- 12 on which neither Simon nor Garfunkel sings



- 1. How many of her CDs feature only Paul Simon? 3
- 2. How many of her CDs feature only Art Garfunkel? 2
- 3. How many feature at least one of these two artists?

- 5 on which both Simon and Garfunkel sing
- 8 total on which Simon sings
- 7 total on which Garfunkel sings
- 12 on which neither Simon nor Garfunkel sings



- 1. How many of her CDs feature only Paul Simon? 3
- 2. How many of her CDs feature only Art Garfunkel? 2
- 3. How many feature at least one of these two artists? 10

There is the cliche that Country–Western songs emphasize three basic themes: love, prison, and trucks. A survey of the local Country–Western radio station produced the following data of songs about:

- 12 truck drivers in love while in prison
- 13 prisoners in love
- 28 people in love
- 18 truck drivers in love
- 3 truck drivers in prison who are not in love
- 2 prisoners not in love and not driving trucks
- 8 people who are out of prison, are not in love, and do not drive trucks
- 16 truck drivers who are not in prison

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | | | |
| 13 | | | |
| 28 | | | |
| 18 | | | |
| 3 | | | |
| 2 | | | |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | | | |
| 28 | | | |
| 18 | | | |
| 3 | | | |
| 2 | | | |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | | | |
| 18 | | | |
| 3 | | | |
| 2 | | | |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | Yes | Maybe | Maybe |
| 18 | | | |
| 3 | | | |
| 2 | | | |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | Yes | Maybe | Maybe |
| 18 | Yes | Maybe | Yes |
| 3 | | | |
| 2 | | | |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | Yes | Maybe | Maybe |
| 18 | Yes | Maybe | Yes |
| 3 | No | Yes | Yes |
| 2 | | | |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | Yes | Maybe | Maybe |
| 18 | Yes | Maybe | Yes |
| 3 | No | Yes | Yes |
| 2 | No | Yes | No |
| 8 | | | |
| 16 | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | Yes | Maybe | Maybe |
| 18 | Yes | Maybe | Yes |
| 3 | No | Yes | Yes |
| 2 | No | Yes | No |
| 8 | No | No | No |
| 16 | | | |
| | | | |

| Number | Love? | Prison? | Trucks? |
|--------|-------|---------|---------|
| 12 | Yes | Yes | Yes |
| 13 | Yes | Yes | Maybe |
| 28 | Yes | Maybe | Maybe |
| 18 | Yes | Maybe | Yes |
| 3 | No | Yes | Yes |
| 2 | No | Yes | No |
| 8 | No | No | No |
| 16 | Maybe | No | Yes |



- 1. Surveyed?
- 2. About truck drivers?
- 3. About prisoners?
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed?
- 2. About truck drivers?
- 3. About prisoners?
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed?
- 2. About truck drivers?
- 3. About prisoners?
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed?
- 2. About truck drivers?
- 3. About prisoners?
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed? 51
- 2. About truck drivers?
- 3. About prisoners?
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed? 51
- 2. About truck drivers? 31
- 3. About prisoners?
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed? 51
- 2. About truck drivers? 31
- 3. About prisoners? 18
- 4. About truck drivers in prison?
- 5. About people not in prison?
- 6. About people not in love?



- 1. Surveyed? 51
- 2. About truck drivers? 31
- 3. About prisoners? 18
- 4. About truck drivers in prison? 15
- 5. About people not in prison?
- 6. About people not in love?


How many songs were. . .

- 1. Surveyed? 51
- 2. About truck drivers? 31
- 3. About prisoners? 18
- 4. About truck drivers in prison? 15
- 5. About people not in prison? 33
- 6. About people not in love?



How many songs were...

- 1. Surveyed? 51
- 2. About truck drivers? 31
- 3. About prisoners? 18
- 4. About truck drivers in prison? 15
- 5. About people not in prison? 33
- 6. About people not in love? 23

Jim Donahue was a section chief for an electric utility company. The employees in his section cut down tall trees (T), climbed poles (P), and spliced wire (W). Donahue submitted the following report to the his manager:

$$n(T) = 45 \qquad n(P \cap W) = 20$$

$$n(P) = 50 \qquad n(T \cap W) = 25$$

$$n(W) = 57 \qquad n(T \cap P \cap W) = 11$$

$$n(T \cap P) = 28 \qquad n(T' \cap P' \cap W') = 9$$

Donahue also stated that 100 employees were included in the report. Why did management reassign him to a new section?











Totals to 99 not 100

Jim Donahue was reassigned to the home economics department of the electric utility company. he interviewed 140 people in a suburban shopping center to find out some of their cooking habits. He obtained the following results. There is a job opening in Siberia. Should he be reassigned yet again?

- 58 use microwave ovens
- 63 use electric ranges
- 58 use gas ranges
- 19 use microwave ovens and electric ranges
- 17 use microwave ovens and gas ranges
- 4 use both gas and electric ranges
- 1 uses all three
- 2 cook only with solar energy











Totals to 142 not 140

Julie Ward, who sells college textbooks, interviewed freshmen on a community college campus to determine what is important to today's students. She found that Wealth, Family, and Expertise topped the list. Her findings can be summarized as:

$$n(W) = 160$$
 $n(E \cap F) = 90$

$$n(F) = 140 \qquad n(W \cap F \cap E) = 80$$

$$n(E) = 130$$
 $n(E') = 95$

$$n(W \cap F) = 95$$
 $n[(W \cup F \cup E)'] = 10$







Counting Principles



We have three choices from A to B and two choices from B to C.

How many ways are there to get from A to C through B?

Counting Principles



We have three choices from A to B and two choices from B to C.

How many ways are there to get from A to C through B? 3 * 2 = 6 ways

If k operations (events, actions,...) are performed in succession where:

Operation 1 can be done in n_1 ways Operation 2 can be done in n_2 ways

Operation k can be done in n_k ways

then the total number of ways the k operations can all be performed is:

```
n_1 * n_2 * n_3 * \cdots * n_k
```

In other words, if you have several actions to do and you must do them all you multiply the number of choices to find the total number of choices. How many outcomes can there be from three flips of a coin?

Action 1:Flip a coinAction 2:Flip a coinAction 3:Flip a coinTotal

How many outcomes can there be from three flips of a coin?

| Action 1: | Flip a coin | $2~{ m ways}$ |
|-----------|-------------|-----------------|
| Action 2: | Flip a coin | $2 { m ways}$ |
| Action 3: | Flip a coin | $2 {\rm ways}$ |
| Total | | |

How many outcomes can there be from three flips of a coin?

| Action 1: | Flip a coin | 2 ways |
|-----------|-------------|----------------------------|
| Action 2: | Flip a coin | 2 ways |
| Action 3: | Flip a coin | 2 ways |
| Total | | $2 * 2 * 2 = 2^3 = 8$ ways |

How many ways are there to form a three letter sequence from the letters in $\{A,B,C,\ldots,Z\}?$

Action 1:Pick a letterAction 2:Pick a letterAction 3:Pick a letter

Total

How many ways are there to form a three letter sequence from the letters in $\{A, B, C, \ldots, Z\}$?

| Action 1: | Pick a letter | $26~\mathrm{ways}$ |
|-----------|---------------|----------------------|
| Action 2: | Pick a letter | $26 \mathrm{~ways}$ |
| Action 3: | Pick a letter | $26 \mathrm{\ ways}$ |

Total

How many ways are there to form a three letter sequence from the letters in $\{A, B, C, \ldots, Z\}$?

| Action 1: | Pick a letter | 26 ways |
|-----------|---------------|------------------------------------|
| Action 2: | Pick a letter | 26 ways |
| Action 3: | Pick a letter | 26 ways |
| Total | | $26 * 26 * 26 = 26^3 = 17576$ ways |

Action 1: Pick a letter Action 2: Pick an unused letter Action 3: Pick an unused letter

Total

| Action 1: | Pick a letter | 26 ways |
|-----------|-----------------------|---------|
| Action 2: | Pick an unused letter | |
| Action 3: | Pick an unused letter | |
| Total | | |

| - I | | |
|-----------|-----------------------|----------------------|
| Action 3: | Pick an unused letter | |
| Action 2: | Pick an unused letter | $25 \ \mathrm{ways}$ |
| Action 1: | Pick a letter | $26 { m ways}$ |

Total

| Action 1: | Pick a letter | $26~{ m ways}$ |
|-----------|-----------------------|----------------------|
| Action 2: | Pick an unused letter | $25 \mathrm{\ ways}$ |
| Action 3: | Pick an unused letter | $24 \mathrm{\ ways}$ |
| | | |

Total

| Action 1: | Pick a letter | 26 ways |
|-----------|-----------------------|---------------------------|
| Action 2: | Pick an unused letter | $25 \mathrm{ways}$ |
| Action 3: | Pick an unused letter | 24 ways |
| Total | | 26 * 25 * 24 = 15600 ways |

Action 1: Pick 3 letters Action 2: Pick 3 digits Total

Action 1:Pick 3 letters 26^3 waysAction 2:Pick 3 digitsTotal

| Action 1: | Pick 3 letters | 26^3 ways |
|-----------|----------------|------------------|
| Action 2: | Pick 3 digits | $10^3~{ m ways}$ |
| Total | | |

| Action 1: | Pick 3 letters | 26^3 ways |
|-----------|----------------|---------------------------------|
| Action 2: | Pick 3 digits | 10^3 ways |
| Total | | $26^3 * 10^3 = 17,576,000$ ways |
Counting with Trees

Tree diagrams consist of nodes (the circles) and branches that connect some nodes.



The nodes represent the possible "states" of a situation.

Branches are the ways or "choices" we have to move to another state.

The "top" node is called the root and it represents the starting state.

Leaves, nodes with no other nodes under them, represent an ending state.

This leads to the following technique:

- Use a tree diagram to illustrate a situation
- Count the number of leaves to find the number of possible outcomes

How many outcomes can there be from three flips of a coin?



How many ways can a best of three Chess series end. Must have a majority to be declared victor.



21 outcomes

If $A \cap B = \emptyset$, then $n(A \cup B) =$

If we are to perform **one** of k operations (events, actions,...) where:

Operation 1 can be done in n_1 ways Operation 2 can be done in n_2 ways \vdots Operation k can be done in n_k ways

then the total number of choices is:

$$n_1 + n_2 + n_3 + \dots + n_k$$

In other words, if you have several actions to do and you are only going to do one of them you add the number of choices to find the total number of choices. If $A \cap B = \emptyset$, then $n(A \cup B) = \mathbf{n}(\mathbf{A}) + \mathbf{n}(\mathbf{B})$.

If we are to perform **one** of k operations (events, actions,...) where:

Operation 1 can be done in n_1 ways Operation 2 can be done in n_2 ways \vdots Operation k can be done in n_k ways

then the total number of choices is:

$$n_1 + n_2 + n_3 + \dots + n_k$$

In other words, if you have several actions to do and you are only going to do one of them you add the number of choices to find the total number of choices. You are hungry and want to order a combo meal from either Taco Hut or Burger Lord. Taco Hut has 6 different combo meals and Burger Lord has 9. How many choices to you have?

Action 1: Order a combo from Taco Hut Action 2: Order a combo from Burger Lord Total You are hungry and want to order a combo meal from either Taco Hut or Burger Lord. Taco Hut has 6 different combo meals and Burger Lord has 9. How many choices to you have?

Action 1:Order a combo from Taco Hut6 choicesAction 2:Order a combo from Burger Lord9 choicesTotal

You are hungry and want to order a combo meal from either Taco Hut or Burger Lord. Taco Hut has 6 different combo meals and Burger Lord has 9. How many choices to you have?

| Action 1: | Order a combo from Taco Hut | 6 choices |
|-----------|--------------------------------|----------------|
| Action 2: | Order a combo from Burger Lord | 9 choices |
| Total | | 6+9=15 choices |

You are hungry and want to order a pizza from either Pizza Place or Pizza Hog. Pizza Place has 6 different toppings and Pizza Hog has 9. Topping can either be on or off of a pizza. How many choices to you have?

Action 1: Order a pizza from Pizza Place Action 2: Order a pizza from Pizza Hog

Total

You are hungry and want to order a pizza from either Pizza Place or Pizza Hog. Pizza Place has 6 different toppings and Pizza Hog has 9. Topping can either be on or off of a pizza. How many choices to you have?

You are hungry and want to order a pizza from either Pizza Place or Pizza Hog. Pizza Place has 6 different toppings and Pizza Hog has 9. Topping can either be on or off of a pizza. How many choices to you have?

| Action 1: | Order a pizza from Pizza Place | $2^6 = 64$ choices |
|-----------|--------------------------------|------------------------|
| Action 2: | Order a pizza from Pizza Hog | $2^9 = 512$ choices |
| Total | | 64 + 512 = 576 choices |